

CHECKMATE[®] OLR-F

A BIOCHEMICAL FOR MATING DISRUPTION OF OMNIVOROUS LEAFROLLER (*Platynota stultana*)

ACTIVE INGREDIENTS:

(E) - 11 - Tetradecen-1-yl acetate	20.57 %
(Z) - 11 - Tetradecen-1-yl acetate	3.19 %
OTHER INGREDIENTS	76.24 %
TOTAL	100.00 %

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Harmful if inhaled. Avoid breathing vapor or spray mist.

User Safety Requirements: Applicators and other handlers must wear long-sleeved shirt and long pants, waterproof gloves, and shoes plus socks. For overhead exposure wear chemical resistant headgear. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

FIRST AID STATEMENT

IF SWALLOWED: Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Center or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a Poison Control Center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a Poison Control Center or doctor for further treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice.

HOT LINE NUMBER: Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment. You may also contact 1-866-326-6737 for emergency medical treatment information.

ENVIRONMENTAL HAZARD

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposing of equipment washwaters.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool place until used. Protect from freezing.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Do not contaminate water when disposing of equipment wash water.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For requirements specific to your state or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS - Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE). The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during application of this product. A restricted entry interval of 0 hours has been established.

METHOD OF APPLICATION: To disrupt mating of the omnivorous leafroller on peaches, nectarines, apricots, plums, prunes, cherries and other stone fruits; apples, quinces, and other pome fruits; almonds and other tree nut crops; bell peppers, chili peppers, cooking peppers and other fruiting vegetables; kiwis; pomegranates; and grapes, wine grapes, and raisins apply **CheckMate[®] OLR-F** at 0.65 fluid ounces (19.22 ml) product per acre [4.47 g a.i./acre] to 1.46 fluid ounces (43.2 ml) product per acre [10 g a.i./acre]. Monitor for insect infestation with traps and by visual inspection of trees and fruit. Do not exceed 22 fluid ounces (644 ml) product/acre/year (150 g a.i./acre/year). Reapply as needed, based on insect monitoring and field scouting programs.

SHAKE WELL BEFORE USING.

DO NOT use **CheckMate[®] OLR-F** from damaged, punctured, or unsealed containers. Use piston, diaphragm, or centrifugal pumps ONLY. DO NOT use roller or gear pumps as they will damage the **CheckMate[®] OLR-F** controlled release system. For best results, apply immediately after mixing. Do not let stand prior to application without thoroughly mixing again.

Ground Equipment: Apply **CheckMate[®] OLR-F** in sufficient water for uniform coverage. Do not apply to point of runoff. Apply to foliage directly. **Aerial Equipment:** Apply **CheckMate[®] OLR-F** in sufficient water for uniform coverage.



RECOMMENDED APPLICATION: For best results, apply at the time or shortly before omnivorous leafroller moths begin to emerge in the spring (biofix), or when the moths are first detected in pheromone traps such as **BioLure[®]**. This product only affects male moths and will have no effect on female moths, eggs or larvae. If application occurs following biofix or during the growing season, the orchard or vineyard must be treated with insecticide treatments that will effectively control hatching larvae until egg laying by previously mated female moths has ceased to occur. In moderate to high insect pressure situations, supplemental insecticide applications during the season may be necessary to provide adequate protection to developing fruit. These supplemental insecticide applications should be made based on trap monitoring, field scouting, and appropriate degree-day models for the local growing area and must be timed to control emerging larvae.

WARRANTY: **Suterra, LLC** warrants that this material conforms to the chemical description on the label. **Suterra** neither makes nor authorizes any agent or representative to make any other warranty of fitness or of merchantability, guarantee or representation, expressed or implied, concerning this material. **Suterra's** maximum liability for breach of this warranty shall not exceed the purchase price of this product. Buyer and user acknowledge and assume all risks and liabilities resulting from the handling, storage, and use of this material.

MANUFACTURED BY:

SUTERRA[™]

Suterra, LLC
213 SW Columbia St.
Bend, OR, U.S.A. 97702-1013
Phone (866) 326-6737
Fax (541) 388-3705

Net Contents: 13.1 fl. oz. (387 ml)
Contains: 6.88 g/a.i./fl. ounce (0.23 g/a.i./ml)
EPA Reg. No. 56336-25 / EPA Est. No. 56336-OR-1
olr#7shpr 10/2003

Lot Number: